

Four-level quantum teleportation, swapping and collective translations of multipartite quantum entanglement

Zai-Zhe Zhong

Department of Physics, Liaoning Normal University, Dalian 116029,
Liaoning, China. E-mail: zhongzaizhecn@yahoo.com.cn

February 1, 2008

Abstract

In this paper, an optimal scheme of four-level quantum teleportation and swapping of quantum entanglement is given. We construct a complete orthogonal basis of the bipartite ququadrit systems. Using this basis, the four-level quantum teleportation and swapping can be achieved according to the standard steps. In addition, associate the above bases with the unextendible product bases and the exact entanglement bases, we prove that in the $2 \times 2 \times 2$ systems or 3×3 systems the collective translations of multipartite quantum entanglement can be realized.

PACC numbers: 03.67.Mn, 03.65.Ud, 03.67.Hk.

Keywords: Ququadrit systems, Bases, Four-level teleportation, Swapping, Collective translations.

Quantum teleportation and swapping are quite interesting and important topics in modern quantum mechanics and quantum information. Following the BBCJPW scheme[1], there have been very many related works (e.g. see the references in [2]). About the swapping of quantum entanglement, the original work was by ZZHE[3]. About the problems how to extend the schemes of quantum teleportation and swapping to the multipartite d -level ($d \geq 3$) cases, there are many works (e.g. see [4-9], and some recent works

related to teleporting states of dimension higher than two, see [10,11]). In this paper, we point out that in the case of ququadrut systems we can find a simple way to achieve quantum teleportation and swapping. We give a new complete orthogonal basis. Using this basis and according to the standard steps, the four-level quantum teleportation and swapping can be realized. The mathematical form of the schemes is simple.

In addition, there is yet another interesting result that we can achieve collective quantum teleportation and collective swapping of quantum entanglement of two or three particles. In this paper, as an application of the above schemes, we prove that this is possible, i.e. we associate the above basis with the unextendible product bases (UPB)[12-16] and the exact entanglement bases (EEB)[17], so that in the $2 \times 2 \times 2$ systems or 3×3 systems collective translations of multipartite quantum entanglement can be realized.

If $H^{(1)}$ and $H^{(2)}$ are the Hilbert spaces of ququadrut states with the natural bases $\{|i\rangle\}$ and $\{|j\rangle\}$ ($i, j = 0, 1, 2, 3$), respectively, then the natural basis of $H = H^{(1)} \otimes H^{(2)}$ is $\{|i\rangle | j\rangle\}$. In the first place, we define a new complete orthogonal basis $\{|W_i\rangle, |X_i\rangle, |Y_i\rangle, |Z_i\rangle\}$ ($i = 0, 1, 2, 3$) of H by

$$\begin{aligned} |W_i\rangle &= \frac{1}{2} (|i\rangle | 0\rangle + |i+1 \bmod 4\rangle | 1\rangle + |i+2 \bmod 4\rangle | 2\rangle + |i+3 \bmod 4\rangle | 3\rangle) \\ |X_i\rangle &= \frac{1}{2} (|i\rangle | 0\rangle + |i+1 \bmod 4\rangle | 1\rangle - |i+2 \bmod 4\rangle | 2\rangle - |i+3 \bmod 4\rangle | 3\rangle) \\ |Y_i\rangle &= \frac{1}{2} (|i\rangle | 0\rangle - |i+1 \bmod 4\rangle | 1\rangle + |i+2 \bmod 4\rangle | 2\rangle - |i+3 \bmod 4\rangle | 3\rangle) \\ |Z_i\rangle &= \frac{1}{2} (|i\rangle | 0\rangle - |i+1 \bmod 4\rangle | 1\rangle - |i+2 \bmod 4\rangle | 2\rangle + |i+3 \bmod 4\rangle | 3\rangle) \end{aligned}$$

Obviously, $\{|W_i\rangle, |X_i\rangle, |Y_i\rangle, |Z_i\rangle\}$ ($i = 0, 1, 2, 3$) are entangled states, they form a complete orthogonal basis of H . The transformation relation between the above basis and the natural basis is

$$\begin{aligned} |i\rangle | 0\rangle &= \frac{1}{2} (|W_i\rangle + |X_i\rangle + |Y_i\rangle + |Z_i\rangle) \\ |i+1 \bmod 4\rangle | 1\rangle &= \frac{1}{2} (|W_i\rangle + |X_i\rangle - |Y_i\rangle - |Z_i\rangle) \\ |i+2 \bmod 4\rangle | 2\rangle &= \frac{1}{2} (|W_i\rangle - |X_i\rangle + |Y_i\rangle - |Z_i\rangle) \\ |i+3 \bmod 4\rangle | 3\rangle &= \frac{1}{2} (|W_i\rangle - |X_i\rangle - |Y_i\rangle + |Z_i\rangle) \end{aligned} \tag{2}$$

Following the standard steps of quantum teleportation, we suppose that Alice holds particle 1 which is in an unknown pure-state $|\phi^{(1)}\rangle = \alpha |0_1\rangle + \beta |1_1\rangle + \gamma |2_1\rangle + \delta |3_1\rangle$, Clara is in a remote place from Alice. Bob holds two particles 2 and 3 which are in some basic state, for instance, in $|X_1^{(2,3)}\rangle = \frac{1}{2}(|1_2\rangle |0_3\rangle + |2_2\rangle |1_3\rangle - |3_2\rangle |2_3\rangle - |0_2\rangle |3_3\rangle)$. The total state is

$$|\Psi_{total}\rangle = |\phi^{(1)}\rangle |X_1^{(2,3)}\rangle = \frac{1}{2}(\alpha |0_1\rangle |1_2\rangle |0_3\rangle + \alpha |0_1\rangle |2_2\rangle |1_3\rangle - \alpha |0_1\rangle |3_2\rangle |2_3\rangle - \alpha |0_1\rangle |0_2\rangle |3_3\rangle + \beta |1_1\rangle |1_2\rangle |0_3\rangle + \dots - \delta |3_1\rangle |0_2\rangle |3_3\rangle) \quad (3)$$

According to Eq.(2), every $|i_1\rangle |j_2\rangle$ can always be expressed by $|W_k^{(1,2)}\rangle$, $|X_k^{(1,2)}\rangle$, $|Y_k^{(1,2)}\rangle$ and $|Z_k^{(1,2)}\rangle$. Substitute them into Eq.(3) and reorganize, the final result is

$$|\Psi_{total}\rangle = \sum_{i=0}^3 \left(|W_i^{(1,2)}\rangle |\phi_{W_i}^{(3)}\rangle + |X_i^{(1,2)}\rangle |\phi_{X_i}^{(3)}\rangle + |Y_i^{(1,2)}\rangle |\phi_{Y_i}^{(3)}\rangle + |Z_i^{(1,2)}\rangle |\phi_{Z_i}^{(3)}\rangle \right) \\ \equiv \frac{1}{4} \left\{ \begin{aligned} & |W_0^{(1,2)}\rangle (\beta |0_3\rangle + \gamma |1_3\rangle - \delta |2_3\rangle - \alpha |3_3\rangle) \\ & + |W_1^{(1,2)}\rangle (\gamma |0_3\rangle + \delta |1_3\rangle - \alpha |2_3\rangle - \beta |3_3\rangle) \\ & + |W_2^{(1,2)}\rangle (\delta |0_3\rangle + \alpha |1_3\rangle - \beta |2_3\rangle - \gamma |3_3\rangle) \\ & + |W_3^{(1,2)}\rangle (\alpha |0_3\rangle + \beta |1_3\rangle - \gamma |2_3\rangle - \delta |3_3\rangle) \\ & + |X_0^{(1,2)}\rangle (\beta |0_3\rangle - \gamma |1_3\rangle - \delta |2_3\rangle - \alpha |3_3\rangle) \\ & + |X_1^{(1,2)}\rangle (\gamma |0_3\rangle - \delta |1_3\rangle + \alpha |2_3\rangle - \beta |3_3\rangle) \\ & + |X_2^{(1,2)}\rangle (\delta |0_3\rangle - \alpha |1_3\rangle + \beta |2_3\rangle - \gamma |3_3\rangle) \\ & + |X_3^{(1,2)}\rangle (-\alpha |0_3\rangle - \beta |1_3\rangle - \gamma |2_3\rangle - \delta |3_3\rangle) \\ & + |Y_0^{(1,2)}\rangle (-\beta |0_3\rangle + \gamma |1_3\rangle + \delta |2_3\rangle - \alpha |3_3\rangle) \\ & + |Y_1^{(1,2)}\rangle (-\gamma |0_3\rangle + \delta |1_3\rangle + \alpha |2_3\rangle - \beta |3_3\rangle) \\ & + |Y_2^{(1,2)}\rangle (-\delta |0_3\rangle - \alpha |1_3\rangle - \beta |2_3\rangle - \gamma |3_3\rangle) \\ & + |Y_3^{(1,2)}\rangle (-\alpha |0_3\rangle + \beta |1_3\rangle + \gamma |2_3\rangle - \delta |3_3\rangle) \\ & + |Z_0^{(1,2)}\rangle (-\beta |0_3\rangle - \gamma |1_3\rangle - \delta |2_3\rangle - \alpha |3_3\rangle) \\ & + |Z_1^{(1,2)}\rangle (-\gamma |0_3\rangle - \delta |1_3\rangle - \alpha |2_3\rangle - \beta |3_3\rangle) \\ & + |Z_2^{(1,2)}\rangle (-\delta |0_3\rangle + \alpha |1_3\rangle + \beta |2_3\rangle - \gamma |3_3\rangle) \\ & + |Z_3^{(1,2)}\rangle (\alpha |0_3\rangle - \beta |1_3\rangle - \gamma |2_3\rangle - \delta |3_3\rangle) \end{aligned} \right\} \quad (4)$$

Now, Bob sends particles 2 and 3 to Alice and Clara respectively, and Alice makes a joint measurement of particles 1, 2. She will obtain of the 16 basic states $\left\{ |W_i^{(1,2)}\rangle, |X_i^{(1,2)}\rangle, |Y_i^{(1,2)}\rangle, |Z_i^{(1,2)}\rangle \right\}$ ($i = 0, 1, 2, 3$) with probability $\frac{1}{16}$ (here we assume that there is such instrument which can distinguish these bases). Simultaneously Clara must obtain a corresponding one state of $\left\{ |\phi_{W_i}^{(3)}\rangle, |\phi_{X_i}^{(3)}\rangle, |\phi_{Y_i}^{(3)}\rangle, |\phi_{Z_i}^{(3)}\rangle \right\}$. We define sixteen unitary matrices as

$$\begin{aligned} U_{W_0} &= \begin{bmatrix} 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{bmatrix}, U_{W_1} = \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} \\ U_{W_2} &= \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \end{bmatrix}, U_{W_3} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} \end{aligned} \quad (5)$$

$$\begin{aligned} U_{X_0} &= \begin{bmatrix} 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{bmatrix}, U_{X_1} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix} \\ U_{X_2} &= \begin{bmatrix} 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \end{bmatrix}, U_{X_3} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} \end{aligned} \quad (6)$$

$$\begin{aligned} U_{Y_0} &= \begin{bmatrix} 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}, U_{Y_1} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} \end{aligned} \quad (7)$$

$$\begin{aligned} U_{Y_2} &= \begin{bmatrix} 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \end{bmatrix}, U_{Y_3} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} \end{aligned} \quad (8)$$

$$\begin{aligned}
U_{Z_0} &= \begin{bmatrix} 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{bmatrix}, \quad U_{Z_1} = \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix} \\
U_{Z_2} &= \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \end{bmatrix}, \quad U_{Z_3} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}
\end{aligned}$$

When Alice informs Clara of her measurement result μ_j , by the classical communications, then Clara at once knows the correct result should be $|\phi^3\rangle = U_{\mu_j} |\phi_{\mu_j}^3\rangle$. Thus we achieve a four-level quantum teleportation.

By using of the above basis, we can also carry out the four-level swapping. For instance, we suppose that Alice holds particle 1, Bob holds particles 2, 3, Clara holds particle 4, particles 1 and 2 are in the entangled state $|X_1^{(1,2)}\rangle$, and particles 3 and 4 are in the entangled state $|X_1^{(3,4)}\rangle$. Therefore the total state is

$$\begin{aligned}
|\Psi_{1234}\rangle &= |X_2^{(1,2)}\rangle |X_2^{(3,4)}\rangle = \frac{1}{4} (|1_1\rangle |0_2\rangle + |2_1\rangle |1_2\rangle - |3_1\rangle |2_2\rangle - |0_1\rangle |3_2\rangle) \\
&\quad \otimes (|1_3\rangle |0_4\rangle + |2_3\rangle |1_4\rangle - |3_3\rangle |2_4\rangle - |0_3\rangle |3_4\rangle) \\
&= \frac{1}{4} (|1_1\rangle |0_2\rangle |1_3\rangle |0_4\rangle + |1_1\rangle |0_2\rangle |2_3\rangle |1_4\rangle - \cdots + |0_1\rangle |3_2\rangle |0_3\rangle |3_4\rangle) \\
&= \frac{1}{8} \left\{ \begin{aligned} &|1_1\rangle \left(|W_3^{(2,3)}\rangle - |X_3^{(2,3)}\rangle - |Y_3^{(2,3)}\rangle + |Z_3^{(2,3)}\rangle \right) |0_4\rangle \\ &+ |1_1\rangle \left(|W_2^{(2,3)}\rangle - |X_2^{(2,3)}\rangle - |Y_2^{(2,3)}\rangle + |Z_2^{(2,3)}\rangle \right) |1_4\rangle \\ &- \cdots + |0_1\rangle \left(|W_3^{(2,3)}\rangle + |X_3^{(2,3)}\rangle + |Y_3^{(2,3)}\rangle + |Z_3^{(2,3)}\rangle \right) |3_4\rangle \end{aligned} \right\} \quad (9)
\end{aligned}$$

For $|1_1\rangle |0_4\rangle, |1_1\rangle |1_4\rangle, \cdots, |0_1\rangle |3_4\rangle$ we use Eq.(2), and we rewrite $|\Psi_{1234}\rangle$

in $(H_2 \otimes H_3) \otimes (H_1 \otimes H_4)$, to obtain

$$|\Psi_{1234}\rangle = \frac{1}{4} \left\{ \begin{array}{l} |W_0^{(2,3)}\rangle |Z_2^{(1,4)}\rangle - |W_1^{(2,3)}\rangle |X_3^{(1,4)}\rangle \\ - |W_2^{(2,3)}\rangle |Y_0^{(1,4)}\rangle + |W_3^{(2,3)}\rangle |W_1^{(1,4)}\rangle \\ + |X_0^{(2,3)}\rangle |X_2^{(1,4)}\rangle - |X_1^{(2,3)}\rangle |X_3^{(1,4)}\rangle \\ - |X_2^{(2,3)}\rangle |X_0^{(1,4)}\rangle - |X_3^{(2,3)}\rangle |X_1^{(1,4)}\rangle \\ - |Y_0^{(2,3)}\rangle |W_2^{(1,4)}\rangle + |Y_1^{(2,3)}\rangle |Y_3^{(1,4)}\rangle \\ + |Y_2^{(2,3)}\rangle |Z_0^{(1,4)}\rangle - |Y_3^{(2,3)}\rangle |Z_1^{(1,4)}\rangle \\ - |Z_0^{(2,3)}\rangle |Y_2^{(1,4)}\rangle - |Z_1^{(2,3)}\rangle |Z_3^{(1,4)}\rangle \\ + |Z_2^{(2,3)}\rangle |W_0^{(1,4)}\rangle + |Z_3^{(2,3)}\rangle |Z_1^{(1,4)}\rangle \end{array} \right\} \quad (10)$$

This means that when Bob makes a joint measurement of particles 2 and 3, so that the wave function $|\Psi_{1234}\rangle$ collapses to only one of the above 16 states (say, $|W_1^{(2,3)}\rangle$) with probability $\frac{1}{16}$, then there appears one corresponding entanglement (say, $|X_3^{(1,4)}\rangle$) between particles 1 and 4, etc. To sum up, the four-level quantum entanglement swapping can be realized by this way.

As an application of the above schemes, we prove that there may be some collective translations of multipartite quantum entanglement. Here we need to use the concepts of the UPB[12-14] and the EEB[17]. Consider an M -partite quantum system $H = \otimes_{i=1}^M H_i$, each H_i is d -dimensional, the total dimensionality of H is $N = d^M$. An UPB of H is a product basis $S = \{|\psi_0\rangle, \dots, |\psi_{m-1}\rangle\}$, which spans a subspace H_S in H , and the complementary subspace $H - H_S$ contains no product state. It is known[12,14] that $m \geq M(d-1) + 1$. Following [17], if $T = \{|\varepsilon_0\rangle, |\varepsilon_2\rangle, \dots, |\varepsilon_{n-1}\rangle\}$ ($m+n = d^M$) is a set of entangled pure-states, and $B = S \cup T$ forms an orthogonal complete basis of H , then we call T an EEB. Obviously, an arbitrary linear combination of $|\varepsilon_0\rangle, |\varepsilon_2\rangle, \dots, |\varepsilon_{n-1}\rangle$ still is an entangled state in H . We call the subspace H_{EES} spanned by T the exact entanglement space (EES), in [17] we have proved the existence of H_{EES} . Evidently $H = H_S \oplus H_{EES}$.

Now we take $m = M(d-1) + 1$ and consider the positive integer solutions d of the following equation

$$d^M - m = d^M - M(d-1) - 1 = 4 \quad (11)$$

There are two solutions, i.e. $M = 2, d = 3$ and $M = 3, d = 2$. This means that in a bipartite qutrit system or in a tripartite qubit system, there is a four dimensional EES (for concrete examples, see [17]). Here for each cases,

we denote the EEB by $\{|\varepsilon_0\rangle, |\varepsilon_1\rangle, |\varepsilon_2\rangle, |\varepsilon_3\rangle\}$, which spans the entanglement space H_{EES} .

Now, we consider a special entanglement problem as follows. The Hilbert space $\mathcal{H} = H_{EES}^{(1)} \otimes H_{EES}^{(2)}$ has an orthogonal complete basis $\{|\varepsilon_i^{(1)}\rangle |\varepsilon_j^{(2)}\rangle\}$ ($i, j = 0, 1, 2, 3$). In $\mathcal{H}$ the general form of a pure-state $|\Phi\rangle$ is

$$|\Phi\rangle = \sum_{i,j=0}^3 f_{ij} |\varepsilon_i^{(1)}\rangle |\varepsilon_j^{(2)}\rangle \quad (12)$$

We define a pure-state $|\Phi\rangle \in \mathcal{H}$ to be ‘separable in $\mathcal{H}$ ’ if and only if it can be decomposed as $|\Phi\rangle = |\Phi_1\rangle |\Phi_2\rangle$, $|\Phi_i\rangle \in H_{EES}^{(i)}$ ($i = 1, 2$); Conversely, $|\Phi\rangle$ is called ‘entangled in $\mathcal{H}$ ’. Notice that since $|\varepsilon_i^{(1)}\rangle, |\varepsilon_j^{(2)}\rangle$ both are entangled states of H themselves, the entanglement in the above definition, in fact, is a special entanglement, i.e. ‘entanglement of entanglement’. Let us make the correspondence

$$|\varepsilon_i^{(k)}\rangle \rightleftharpoons |i_k\rangle \quad (i = 0, 1, 2, 3; k = 1, 2) \quad (13)$$

i.e. we regard every basis state $|i_k\rangle$ ($k = 1, 2$) an entangled state in $H_{EES}^{(k)} \subset H$ ($k = 1, 2$), then the entanglement problems of $\mathcal{H}^{(1,2)} = H_{EES}^{(1)} \otimes H_{EES}^{(2)}$ are just a ququadrit entanglement problems, and we can use the basis as $\{ |W_i^{(1,2)}\rangle, |X_i^{(1,2)}\rangle, |Y_i^{(1,2)}\rangle, |Z_i^{(1,2)}\rangle \}$ ($i = 0, 1, 2, 3$), etc..

Now for a tripartite qubit system, suppose that Alice holds three particles 1, 2, 3 which are in an unknown state $|\phi^{(123)}\rangle = \alpha |\varepsilon_0^{(123)}\rangle + \beta |\varepsilon_1^{(123)}\rangle + \gamma |\varepsilon_2^{(123)}\rangle + \delta |\varepsilon_3^{(123)}\rangle$, Bob holds six particles 4, 5, $\dots$, 9 which are in the entangled state $|X_1^{(456,789)}\rangle = \frac{1}{2} \begin{pmatrix} |\varepsilon_1^{(456)}\rangle |\varepsilon_0^{(789)}\rangle + |\varepsilon_2^{(456)}\rangle |\varepsilon_1^{(789)}\rangle \\ - |\varepsilon_3^{(456)}\rangle |\varepsilon_2^{(789)}\rangle - |\varepsilon_0^{(456)}\rangle |\varepsilon_3^{(789)}\rangle \end{pmatrix}$, then, as above, we can complete the quantum teleportation of $|\phi^{(123)}\rangle$ from Alice to remote Clara.

Similarly, suppose that there are twelve spin- $\frac{1}{2}$ particles 1, 2, $\dots$, 6, 1', 2', $\dots$, 6', Alice holds particles 1, 2, 3, Bob holds six particles 4, 5, 6, 1', 2', 3', Clara holds particles 4', 5', 6', particles 1, 2, 3 and 4, 5, 6 are in the entangled state $|X_1^{(123,456)}\rangle$, and particles 1', 2', 3' and 4', 5', 6' are in the entangled state $|X_1^{(1'2'3',4'5'6')}\rangle$. Therefore the total state is $|\Psi_{1234}\rangle = |X_2^{(123,456)}\rangle |X_2^{(1'2'3',4'5'6')}\rangle$, and according to the above steps of swapping, it can be swapped into the entanglement between particle groups (1, 2, 3) and (4', 5', 6'), and between

groups (4, 5, 6) and (1', 2', 3'), etc.. The case of a bipartite qutrit systems is similar.

Obviously, the above quantum teleportation and swapping, in fact, are some collective translations of quantum entanglement. Especially, the foundations of this method are the four-level quantum teleportation, swapping, and EEB.

At last, we briefly mention the problem of generalization. We find that if the above scheme is directly used to the three-level systems, then some information will be lost. Next, the schemes also can be extended to the N-level ($N \geq 5$) systems, however the results are quite complex.

Conclusion: In the ququadrit systems there are special bases. Using these bases, we can simply realize the four-level quantum teleportation and swapping. As an application of the above schemes, it is proved that in the $2 \times 2 \times 2$ or 3×3 systems there may be some collective translations of multipartite quantum entanglement.

References

- [1] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Phys. Rev. Lett., **70**(1993)1895.
- [2] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information*. Cambridge University Press (2000).
- [3] M. Zukowski, A. Zeilinger, M. A. Horne, and A. Ekert, Phys. Rev. Lett., **71**(1993)4287.
- [4] F. Verstraete and H. Verschelde, Phys. Rev. Lett., **90**(2003)097901.
- [5] J. D. Zhou, G. Hou, and Y. D. Zhang, Phys. Rev. A, **64**(2001)012301
- [6] W. Son, J. H. Lee, M. S. Kim, and Y. J. Park, Phys. Rev. A, **64**(2001)064304.
- [7] J. H. Lee, H. Min, and S. D. Oh, Phys. Rev. A, **66**(2002)052318.
- [8] A. Grudka, Acta Phys. Slov., **54**(2004)91.
- [9] A. Grudka and R. W. Chhajlany, Acta Phys. Pol., **A 104**(2003)409.

- [10] G. Rigolin, Phys. Rev. A, **71**(2005)032303.
- [11] X. H. Ge and Y. G. Shen, Phys. Lett. B**606** (2005)184.
- [12] C. H. Bennett , D. P. DiVincenzo , T. Mor, P. W. Shor, J. A. Smolin ,
and B. M. Terhal, Phys. Rev. Lett., **82**(1999)5385.
- [13] B. M. Terhal, Lin. Alg. Appl., **323**(2000)61.
- [14] D. P. DiVincenzo , T. Mor, P. W. Shor, J. A. Smolin and B. M. Terhal,
Comm. Math. Phys., **238**(2003)379.
- [15] S. Chaturvedi, Phys. Rev. A, 65(2002)042322.
- [16] A. O. Pittenger, Linear Algebr. Appl., **359**(2003)235.
- [17] Z. Z. Zhong, Phys. Rev. A, **70**(2004)044302.